Phase II - Arrival of Recovery Team to Withdrawal of Support

Communication and Huddle


Do you perform a huddle with hospital, OPO and transplant teams, including recovery surgeons, prior to withdrawal of support?



Does your OPO, or do any hospitals within your OPO, require separate physicians for end-of-life care and declaration of cardiovascular death?


Field	Count
OPO requires separate physicians	6.38% 3
Hospital requires separate physicians	4.26% 2
Both require separate physicians	8.51% 4
This is not required by either	80.85% 38
Unknown	0.00% -
Total	47

In what proportion of hospitals in your DSA, is withdrawal of care allowed in the following locations:




Field	Mean	Standa	rd Deviation	Variance	Resp	onses
Operating Room	3.84		1.59	2.53		45
PACU/Holding Room	3.29		1.77	3.13		41
ICU only	1.53		1.24	1.54		34
Other	1.00		0.00	0.00		11
Field	0-20%	21-40%	41-60%	61-80%	> 80%	Total
Operating Room	8	3	4	3	27	45
PACU/Holding Room	13	3	2	5	18	41
ICU only	28	1	1	1	3	34
Other	11	0	0	0	0	11

Does location change if:




Field	Yes	No
Liver recovery is anticipated?	9%	23%
Lung recovery is anticipated?	15%	21%
A pediatric patient?	21%	20%
Many family present for extubation?	45%	14%
On ECMO?	11%	22%

Does your OPO's policy allow pre-withdrawal of life support prep/drape?




Do you believe that family presence is a barrier to successful DCD?



Do you offer family members the opportunity to be present for withdrawal of support?

Field	Count
Yes	48
No	-
Total	48

Does your OPO allow the administration of pre-withdrawal medications?



If yes, which are allowed (select all that apply):

